Molecular Brain Research, 14 (1992) 369–372
Elsevier

Author Index

Abe, H., Watanabe, M., Goto, K., Sakagami, H. and Kondo, H., Localization of gene expression of calreticulin in the brain of adult mouse, 337
Abe, T., see Morita, T., 143
Adler, E.M., see Fink, J.S., 186
Adrien, J., see Miquel, M.C., 357
Amano, T., Yamakuni, T., Okabe, N., Kuwahara, R., Ozawa, F. and Hishinuma, F., Regulation of nerve growth factor and nerve growth factor receptor production by NMDA in C6 glioma cells, 35
Amoroso, D., see Giorgi, S., 277
Araki, K., see Morita, T., 143
Augood, S.J., see Herbison, A.E., 310

Baden, D.G., see Edwards, R.A., 64
Baskys, A., see Billia, F., 101
Bassetti, M.A., see Zucconi, G.G., 1
Beck, J., see Bialy, M., 352
Bialy, M., Nikolaev, E., Beck, J. and Kaczmarek, L., Delayed c-*fos* expression in sensory cortex following sexual learning in male rats, 352
Billia, F., Baskys, A., Carlen, P.L. and De Boni, U., Rearrangement of centromeric satellite DNA in hippocampal neurons exhibiting long-term potentiation, 101
Bock, S., see Lin, L.-H., 147
Boissin, J., see Boissin-Agasse, L., 57
Boissin-Agasse, L., De Bouard, V., Roch, G. and Boissin, J., In situ hybridization of GnRH mRNA in the rat and the mink hypothalamus using biotinylated synthetic oligonucleotide probes, 57
Bono, C., see Eva, C., 344
Borrelli, E., see Micheletti, G., 363
Bowman, C.L., see Edelman, A.M., 27
Branch, A.D., Unterwald, E.M., Lee, S.E. and Kreek, M.J., Quantitation of preproenkephalin mRNA levels in brain regions from male Fischer rats following chronic cocaine treatment using a recently developed solution hybridization assay, 231
Bravo, R., see Herdegen, T., 155
Brené, S., Lindefors, N. and Persson, H., Midbrain dopamine neurons regulate preprotachykinin-A mRNA expression in the rat forebrain during development, 13
Buck, K.J., see Zahniser, N.R., 196
Bunel, D.T., see Valentijn, K., 221
Byrnes, A.P., see McClintock, T.S., 273

Carlen, P.L., see Billia, F., 101
Carlsson, B., see Eriksson, P.S., 317
Carpenter, K., see Lin, L.-H., 147
Carsana, A., see Zucconi, G.G., 1
Chen, K., Yang, W., Grimsby, J. and Shih, J.C., The human 5-HT_2 receptor is encoded by a multiple intron–exon gene, 20
Cho-Lee, J., see Edelman, A.M., 27
Chrisman, C.L., see Waldbieser, G.C., 87
Consolo, S., see Giorgi, S., 277
Cortner, J., see Mack, K.J., 117
Cosi, C., see Zucconi, G.G., 1
Cunningham, A., see Dowsing, B.J., 250
Curella, P., see Zahniser, N.R., 196

Dahl, S.G., see Edvardsen, , 166
De Boni, U., see Billia, F., 101
De Bouard, V., see Boissin-Agasse, L., 57
Dicou, E., Nerve growth factor precursors in the rat thyroid and hippocampus, 136
Dixon, J.E., see Waldbieser, G.C., 87
Doucet, E., see Miquel, M.C., 357
Dowsing, B.J., Gooley, A.A., Gunning, P., Cunningham, A. and Jeffrey, P.L., Molecular cloning and primary structure of the avian Thy-1 glycoprotein, 250
Dragunow, M., Sirimanne, E., Lawlor, P.A., Williams, C. and Gluckman, P., Accumulation of calcitonin-gene related peptide-like immunoreactivity after hypoxic-ischaemic brain injury in the infant rat, 267
Dyer, D., Tigyi, G. and Miledi, R., The effect of active serum albumin on PC12 cells: I. Neurite retraction and activation of the phosphoinositide second messenger system, 293
Dyer, D., Tigyi, G. and Miledi, R., The effect of active serum albumin on PC12 cells: II. Intracellular Ca^{2+} transients and their role in neurite retraction, 302

Edelman, A.M., Higgins, D.M., Bowman, C.L., Haber, S.N., Rabin, R.A. and Cho-Lee, J., Myosin light chain kinase is expressed in neurons and glia: immunoblotting and immunocytochemical studies, 27
Edvardsen, , Sylte, I. and Dahl, S.G., Molecular dynamics of serotonin and ritanserin interacting with the 5-HT_2 receptor, 166
Edwards, E., see Lachman, H.M., 94
Edwards, R.A., Trainer, V.L. and Baden, D.G., Brevetoxins bind to multiple classes of sites in rat brain synaptosomes, 64
Eriksson, P.S., Carlsson, B., Isaksson, O.G.P., Hansson, E. and Rönnbäck, L., Altered amounts of G-protein mRNA and cAMP accumulation after long-term opioid receptor stimulation of neurons in primary culture from the rat cerebral cortex, 317
Eva, C., Fusco, M., Bono, C., Tria, M.A., Gamalero, S.R., Leon, A. and Genazzani, E., Nerve growth factor modulates the expression of muscarinic cholinergic receptor messenger RNA in telencephalic neuronal cultures from newborn rat brain, 344

Farnham, P.J., see Mack, K.J., 117
Fiallos-Estrada, C.E., see Herdegen, T., 155
Fink, J.S., Weaver, D.R., Rivkees, S.A., Peterfreund, R.A., Pollack, A.E., Adler, E.M. and Reppert, S.M., Molecular cloning of the rat A_2 adenosine receptor: selective co-expression with D_2 dopamine receptors in rat striatum, 186
Forloni, G., see Giorgi, S., 277
Fossom, L.H., see Goldstein, M.E., 79
Furia, A., see Zucconi, G.G., 1
Fusco, M., see Eva, C., 344

Gall, C.M., see Yount, G.L., 139
Gallager, D.W., see Primus, R.J., 179
Gamalero, S.R., see Eva, C., 344
Genazzani, E., see Eva, C., 344
Geraerts, W.P.M., see Smit, A.B., 7
Giorgi, S., Forloni, G., Amoroso, D. and Consolo, S., Expression of GAL mRNA in rat hypothalamus: effect of frontal deafferentation and colchicine treatment, 277
Gluckman, P., see Dragunow, M., 267
Goldstein, M.E., Tank, A.W., Fossom, L.H. and Hamill, R.W., Molecular aspects of the regulation of tyrosine hydroxylase by testosterone, 79
Gooley, A.A., see Dowsing, B.J., 250
Goto, K., see Abe, H., 337
Grimsby, J., see Chen, K., 20
Gundlach, A.L., see Rutherfurd, S.D., 261
Gunning, P., see Dowsing, B.J., 250

Haber, S.N., see Edelman, A.M., 27
Haby, C., see Micheletti, G., 363
Hagihara, K., Hirata, S., Osada, T., Hirai, M. and Kato, J., Distribution of cells containing progesterone receptor mRNA in the female rat di- and telencephalon: an in situ hybridization study, 239
Hale, C., see Tümer, N., 51
Hamill, R.W., see Goldstein, M.E., 79
Hamon, M., see Miquel, M.C., 357
Hansson, E., see Eriksson, P.S., 317
Harris, R.A., see Zahniser, N.R., 196
Hartnick, C., see Lachman, H.M., 94
Heidenreich, K.A., see Zahniser, N.R., 196
Henn, F.A., see Lachman, H.M., 94
Herbison, A.E., Augood, S.J. and McGowan, E.M., Expression of glutamic acid decarboxylase messenger RNA in rat medial preoptic area neurones during the oestrous cycle and after ovariectomy, 310
Herdegen, T., Fiallos-Estrada, C.E., Schmid, W., Bravo, R. and Zimmermann, M., The transcription factors c-JUN, JUN D and CREB, but not FOS and KROX-24, are differentially regulated in axotomized neurons following transection of rat sciatic nerve, 155
Higgins, D.M., see Edelman, A.M., 27
Hirai, M., see Hagihara, K., 239
Hirata, S., see Hagihara, K., 239
Hishinuma, F., see Amano, T., 35

Howells, R.D., see Yin, J., 213
Hunt, S.P., see Rusak, B., 124

Iizuka, M., see Satoh, T., 131
Isaksson, O.G.P., see Eriksson, P.S., 317

Jacobs, A.A., see Primus, R.J., 179
Jeffrey, P.L., see Dowsing, B.J., 250
Joosse, J., see Smit, A.B., 7

Kaczmarek, L., see Bialy, M., 352
Kato, J., see Hagihara, K., 239
Keir, W.J., see Zahniser, N.R., 196
Kempf, E., see Micheletti, G., 363
Klein, R.L., see Zahniser, N.R., 196
Kondo, H., see Abe, H., 337
Kreek, M.J., see Branch, A.D., 231
Kushiya, E., see Morita, T., 143
Kuwahara, R., see Amano, T., 35

Lachman, H.M., Papolos, D.F., Weiner, E.D., Ramazankhana, R., Hartnick, C., Edwards, E. and Henn, F.A., Hippocampal neuropeptide Y mRNA is reduced in a strain of learned helpless resistant rats, 94
Landreth, G., see Schanen, N.C., 43
Lannes, B., see Micheletti, G., 363
Lawler, J., see Tümer, N., 51
Lawlor, P.A., see Dragunow, M., 267
Lee, J.A., see Yin, J., 213
Lee, S.E., see Branch, A.D., 231
Leon, A., see Eva, C., 344
Lerner, M.R., see McClintock, T.S., 273
Lin, L.-H., Bock, S., Carpenter, K., Rose, M. and Norden, J.J., ynthesis and transport of GAP-43 in entorhinal cortex neurons and perforant pathway during lesion-induced sprouting and reactive synaptogenesis , 147
Lindefors, N., see Brené, S., 13
Louis, W.J., see Rutherfurd, S.D., 261
Luiten, P.G.M., see Van der Zee, E.A., 326

Mack, K.J., Cortner, J., Mack, P. and Farnham, P.J., *krox* 20 messenger RNA and protein expression in the adult central nervous system, 117
Mack, P., see Mack, K.J., 117
McClintock, T.S., Byrnes, A.P. and Lerner, M.R., Molecular cloning of a G-protein α_i subunit from the lobster olfactory organ, 273
McGeer, P.L., see Walker, D.G., 109
McGowan, E.M., see Herbison, A.E., 310
McQuilkin, S.J., see Zahniser, N.R., 196
McNaughton, L., see Rusak, B., 124
Meester, I., see Smit, A.B., 7
Meguro, H., see Morita, T., 143
Mhatre, M.C. and Ticku, M.K., Aging related alterations in $GABA_A$ receptor subunit mRNA levels in Fischer rats, 71
Micheletti, G., Lannes, B., Haby, C., Borrelli, E., Kempf, E., Warter, J.-M. and Zwiller, J., Chronic administration of NMDA antagonists induces D_2 receptor synthesis in rat striatum, 363
Miledi, R., see Dyer, D., 293
Miledi, R., see Dyer, D., 302
Miller, C.L., see Zahniser, N.R., 196
Minth, C.D., see Waldbieser, G.C., 87
Miquel, M.C., Doucet, E., Riad, M., Adrien, J., Vergé, D. and Hamon, M., Effect of the selective lesion of serotoninergic neurons on the regional distribution of 5-HT_{1A} receptor mRNA in the rat brain, 357
Mishina, M., see Morita, T., 143
Monden, T., see Satoh, T., 131
Mori, M., see Satoh, T., 131
Mori, K.J., see Morita, T., 143
Morita, T., Sakimura, K., Kushiya, E., Yamazaki, M., Meguro, H., Araki, K., Abe, T., Mori, K.J.and Mishina, M., loning and functional expression of a cDNA encoding the mouse $\beta 2$ subunit of the kainate-selective glutamate receptor channel , 143

Nikolaev, E., see Bialy, M., 352
Norden, J.J., see Lin, L.-H., 147

Okabe, N., see Amano, T., 35
Osada, T., see Hagihara, K., 239
Ozawa, F., see Amano, T., 35

Palmieri, M., see Zucconi, G.G., 1
Papolos, D.F., see Lachman, H.M., 94
Persson, H., see Brené, S., 13
Peterfreund, R.A., see Fink, J.S., 186
Pollack, A.E., see Fink, J.S., 186
Prasad, C., see Sakata, M., 282
Primus, R.J., Jacobs, A.A. and Gallager, D.W., Developmental profile of polyadenylated and non-polyadenylated $GABA_A$ receptor subunit mRNAs, 179
Pujol, M.J., see Vendrell, M., 285

Rabin, R.A., see Edelman, A.M., 27
Ramazankhana, R., see Lachman, H.M., 94
Reppert, S.M., see Fink, J.S., 186
Riad, M., see Miquel, M.C., 357
Rivkees, S.A., see Fink, J.S., 186
Robertson, H.A., see Rusak, B., 124
Roch, G., see Boissin-Agasse, L., 57
Rönnbäck, L., see Eriksson, P.S., 317
Rose, M., see Lin, L.-H., 147
Rusak, B., McNaughton, L., Robertson, H.A. and Hunt, S.P., Circadian variation in photic regulation of immediate-early gene mRNAs in rat suprachiasmatic nucleus cells, 124
Rutherfurd, S.D., Widdop, R.E., Louis, W.J. and Gundlach, A.L., Preprogalanin mRNA is increased in vagal motor neurons following axotomy, 261

Sakagami, H., see Abe, H., 337
Sakata, M. and Prasad, C., Transient decrease in rat striatal D_2 dopamine receptor mRNA level after acute haloperidol treatment, 282
Sakimura, K., see Morita, T., 143
Satoh, T., Yamada, M., Monden, T., Iizuka, M. and Mori, M., Cloning of the mouse hypothalamic preprothyrotropin-releasing hormone (TRH) cDNA and tissue distribution of its mRNA, 131
Scarth, B.A., see Shaw, C., 207
Schanen, N.C. and Landreth, G., Isolation and characterization of microtubule-associated protein 2 (MAP2) kinase from rat brain, 43
Schmid, W., see Herdegen, T., 155
Schröder, H., see Van der Zee, E.A., 326
Serratosa, J., see Vendrell, M., 285
Shaw, C. and Scarth, B.A., Age-dependent regulation of $GABA_A$ receptors in neocortex, 207
Shih, J.C., see Chen, K., 20
Sikela, J.M., see Zahniser, N.R., 196
Sirimanne, E., see Dragunow, M., 267
Smit, A.B., Thijsen, S.F.T., Geraerts, W.P.M., Meester, I., Van Heerikhuizen, H. and Joosse, J., Characterization of a cDNA clone encoding molluscan insulin-related peptide V of *Lymnaea stagnalis*, 7
Streefland, C., see Van der Zee, E.A., 326
Strong, R., see Tümer, N., 51
Strosberg, A.D., see Van der Zee, E.A., 326
Sylte, I., see Edvardsen, , 166

Tank, A.W., see Goldstein, M.E., 79
Thijsen, S.F.T., see Smit, A.B., 7
Ticku, M.K., see Mhatre, M.C., 71
Tigyi, G., see Dyer, D., 293
Tigyi, G., see Dyer, D., 302
Trainer, V.L., see Edwards, R.A., 64
Tria, M.A., see Eva, C., 344
Tümer, N., Hale, C., Lawler, J. and Strong, R., Modulation of tyrosine hydroxylase gene expression in the rat adrenal gland by exercise: effects of age, 51
Tusell, J.M., see Vendrell, M., 285

Unterwald, E.M., see Branch, A.D., 231

Valentijn, K., Bunel, D.T. and Vaudry, H., ω-Conotoxin- and nifedipine-insensitive voltage-operated calcium channels mediate K^+-induced release of prothyrotropin-releasing hormone-connecting peptides Ps4 and Ps5 from perifused rat hypothalamic slices, 221
Van der Zee, E.A., Streefland, C., Strosberg, A.D., Schröder, H. and Luiten, P.G.M., Visualization of cholinoceptive neurons in the rat neocortex: colocalization of muscarinic and nicotinic acetylcholine receptors, 326
Van Heerikhuizen, H., see Smit, A.B., 7
Vaudry, H., see Valentijn, K., 221
Vendrell, M., Pujol, M.J., Tusell, J.M. and Serratosa, J., Effect of different convulsants on calmodulin levels and proto-oncogene c-*fos* expression in the central nervous system, 285
Vergé, D., see Miquel, M.C., 357

Waldbieser, G.C., Minth, C.D., Chrisman, C.L. and Dixon, J.E., Tissue-specific expression of the human neuropeptide Y gene in transgenic mice, 87
Walker, D.G. and McGeer, P.L., Complement gene expression in human brain: comparison between normal and Alzheimer disease cases, 109
Warter, J.-M., see Micheletti, G., 363

Watanabe, M., see Abe, H., 337
Weaver, D.R., see Fink, J.S., 186
Weiner, E.D., see Lachman, H.M., 94
White, J.D., see Yount, G.L., 139
Widdop, R.E., see Rutherfurd, S.D., 261
Williams, C., see Dragunow, M., 267
Wilson-Shaw, D., see Zahniser, N.R., 196

Yamada, M., see Satoh, T., 131
Yamakuni, T., see Amano, T., 35
Yamazaki, M., see Morita, T., 143
Yang, W., see Chen, K., 20
Yin, J., Lee, J.A. and Howells, R.D., Stimulation of c-*fos* and c-*jun* gene expression and down-regulation of proenkephalin gene expression in C6 glioma cells by endothelin-1, 213
Yount, G.L., Gall, C.M. and White, J.D., Limbic seizures increase cyclophilin mRNA levels in rat hippocampus, 139

Zahniser, N.R., Buck, K.J., Curella, P., McQuilkin, S.J., Wilson-Shaw, D., Miller, C.L., Klein, R.L., Heidenreich, K.A., Keir, W.J., Sikela, J.M., and Harris, R.A., $GABA_A$ receptor function and regional analysis of subunit mRNAs in long-sleep and short-sleep mouse brain, 196
Zimmermann, M., see Herdegen, T., 155
Zucconi, G.G., Cosi, C., Palmieri, M., Furia, A., Bassetti, M.A. and Carsana, A., A pancreatic-like ribonuclease is synthesized in rat brain, 1
Zwiller, J., see Micheletti, G., 363